

Cannabis Use During Early Pregnancy and Child Diagnoses of Depressive and Anxiety Disorders

Cannabis

2026

© Author(s) 2026

researchmj.org

10.26828/cannabis/2026/000362



Kelly C. Young-Wolff^{1,2,3}, Mahlet Shenkute¹, Nina Oberman¹, Stacey E. Alexeeff¹, Lisa A. Croen^{1,3}, Jamie O. Lo^{4,5}, Deborah Ansley⁶, Carley Castellanos⁶, Kathryn K. Ridout^{1,3,7}, Matthew E. Hirschtritt^{1,2,8}, & Lindsay A. Avalos^{1,3}

¹Division of Research, Kaiser Permanente Northern California

²Department of Psychiatry and Behavioral Sciences, University of California, San Francisco

³Department of Health Systems Science, Bernard J. Tyson Kaiser Permanente School of Medicine

⁴Department of Obstetrics and Gynecology, Oregon Health and Science University

⁵Division of Reproductive and Developmental Science, Oregon National Primate Research Center

⁶The Permanente Medical Group, Regional Offices

⁷Department of Psychiatry, Kaiser Permanente, Santa Rosa Medical Center

⁸Department of Psychiatry, Kaiser Permanente, Oakland Medical Center

ABSTRACT

Objective: Prenatal cannabis use is associated with adverse pregnancy and neonatal outcomes, but research on its association with child anxiety and depressive disorders is limited. This study aimed to test the hypothesis that prenatal cannabis use is associated with child anxiety or depressive disorder diagnoses.

Method: Population-based retrospective birth cohort study of children ($N = 115,553$) born between 1/1/2011–12/31-2017 to pregnant individuals ($N = 97,376$) universally screened for prenatal cannabis use. Cox proportional hazards regression models examined associations between cannabis use during early pregnancy (at ~8-10 weeks gestation based on self-reported use or a positive urine toxicology test for delta-9-tetrahydrocannabinol [THC]) and child anxiety and depressive disorders from ages 6 to 13 years based on diagnosis codes, adjusting for maternal sociodemographic and clinical characteristics. **Results:** The median age at pregnancy onset was 31 (inter quartile range [IQR] = 6) years. Most pregnancies (61.4%) were to non-White individuals; 4.4% screened positive for any cannabis use; 3.7% had a positive toxicology test and 1.9% self-reported use. Overall, 9,230 children were diagnosed with an anxiety disorder at median age of 8 (IQR = 3) years and 1,224 with depressive disorder at median age of 9 (IQR = 3) years. Prenatal cannabis use was associated with a lower risk of child anxiety disorders (aHR:0.84, 95%CI: 0.75-0.95), which remained significant when defined by a toxicology test but not by self-report. Maternal prenatal cannabis use was not associated with child depressive disorders (aHR:1.15, 95%CI: 0.85-1.54). **Conclusions:** Prenatal cannabis use during early pregnancy was not associated with an increased risk of offspring early onset anxiety or depressive disorders.

Key words: = mental health; depression; anxiety; marijuana; cannabis; pregnancy

Corresponding Author: Kelly C. Young-Wolff, PhD, MPH, Division of Research, Kaiser Permanente Northern California, Pleasanton, California 94612. Email: Kelly.c.young-wolff@kp.org

Rates of cannabis use during pregnancy are increasing in the US (Volkow et al., 2019)(Young-Wolff, Chi, et al., 2024), paralleling the growing accessibility, social acceptability, and legalization of cannabis (DISA Global Solutions; Young-Wolff et al., 2022). Despite national guidelines advising against prenatal cannabis use ("Committee Opinion No. 722: marijuana use during pregnancy and lactation," 2017), pregnant individuals report using cannabis to manage morning sickness and mental health symptoms (Bayrampour et al., 2019; Chang et al., 2019; Foti et al., 2023).

Prenatal cannabis use crosses the placenta (Blackard & Tennes, 1984; Lee & Chiang, 1985) and is associated with increased risk of adverse neonatal outcomes, including low birth weight, small-for-gestational-age offspring, and preterm birth (Avalos, Adams, et al., 2024; Baia & Domingues, 2022; Lo et al., 2023). Emerging evidence also suggests an association between prenatal cannabis use and increased risk of adverse maternal outcomes and childhood outcomes, including attention and behavioral difficulties, although results remain mixed (Avalos, Oberman, et al., 2024; Avalos, Shenkute, et al., 2024; Bandoli et al., 2021; Chabarría et al., 2016; Cioffredi et al., 2022; Corsi et al., 2020; Corsi et al., 2019; De Genna et al., 2022; Fried & Watkinson, 1988, 1990; Fried et al., 1992; Garrison-Desany et al., 2022; Grant et al., 2020; Koto et al., 2022; Paul et al., 2021; Paul et al., 2023; Petrangelo et al., 2019; Sharapova et al., 2018; Suján et al., 2022; Tchuente et al., 2022; Torres et al., 2020; Warshak et al., 2015; Young-Wolff, Adams, et al., 2024; Young-Wolff et al., 2025).

The endocannabinoid system (ECS) plays a critical role in healthy brain development and the regulation of emotional responses. ECS receptors (e.g., CB1 and CB2) are involved in neurotransmitter release and neural development. Cannabinoids, including tetrahydrocannabinol (THC) can interact with these receptors, and prior studies suggest that exposure to cannabis in utero may alter ECS and affect neural circuit development involved in stress regulation (Mulligan & Hamre, 2023). Importantly, while considerable research has focused on the relationship between prenatal cannabis exposure and neonatal outcomes or externalizing behaviors and attention-related issues in children, the association between

prenatal cannabis and childhood anxiety and depressive disorders has been overlooked. A recent meta-analysis of initial studies found no association between prenatal cannabis use and child depression (adjusted pooled odds ratio [OR] = 0.72, 95% CI 0.11-4.57) or anxiety (adjusted pooled OR = 1.34, 95% CI 0.79-2.29; (Bassalov et al., 2024). Studies to date have limitations, including the use of older data, self-reported cannabis use, or a combined measure of prenatal and postpartum cannabis use. Furthermore, studies have relied on parent or child self-reported mental health symptoms rather than clinician-documented diagnoses.

Given potential underreporting of prenatal cannabis use (Young-Wolff et al., 2020) and the increasing strength of cannabis products (ElSohly et al., 2016), research that uses both self-report and toxicology testing to assess how prenatal cannabis use relates to clinician-documented diagnoses of child anxiety and depressive disorders is critically needed. In this large, retrospective cohort study, we tested the hypothesis that maternal prenatal cannabis use was associated with the development of early onset of anxiety and depressive disorders in their children born between 2011 and 2017, using data from pregnant individuals universally screened for cannabis use through self-report and urine toxicology testing.

METHODS

Setting and Sample

This study took place in Kaiser Permanente Northern California (KPNC), an integrated healthcare delivery system providing healthcare to >4.6 million patients representative of the insured population in the region. IRB approval was obtained from KPNC and the state of California (governing the use of birth certificate data) with a waiver of consent. STROBE reporting guidelines for cohort studies were followed.

We conducted a retrospective, longitudinal cohort study of pregnant individuals and their singleton children born at a KPNC facility between 1/1/2011–12/31/2017. Eligibility criteria included continuous KPNC enrollment 1-year before pregnancy onset through delivery (allowing for <3-month gaps), attendance at ≥1 KPNC prenatal visit, a response to the question about prenatal

cannabis use, and a prenatal toxicology test for THC. Exclusion criteria included a prenatal prescription fill for a teratogenic, antineoplastic, or antiepileptic drug and missing parity or address data. Children not enrolled in KPNC at age 6 were excluded (eFigure 1). Individuals with >1 pregnancy during the study contributed >1 child to the analysis.

Measures

Data were from the electronic health record (EHR) unless otherwise specified.

Prenatal cannabis use. Pregnant individuals receiving care in KPNC are screened for prenatal cannabis use at entrance to prenatal care (at ~8-10 weeks gestation) via a self-administered questionnaire that assesses frequency of use since pregnancy (“none,” “monthly or less,” “weekly,” and “daily”) and a urine toxicology test to which they consent (Young-Wolff et al., 2017). The primary exposure was any cannabis use during early pregnancy based on self-reported prenatal cannabis use or a positive toxicology test (eAppendix 1). As a secondary exposure, we also examined frequency of prenatal cannabis use (never, monthly or less, weekly, daily, or unknown [no self-reported use and a positive toxicology test]).

Child anxiety and depressive disorders. Anxiety (F40, F40.X, F41, F41.X, F43, F43.X, F94, F94.X) and depressive disorders (F32, F32.X, F33, F33.X) were based ICD-10 diagnostic codes (eAppendix 4). The earliest diagnosis date was used to determine each outcome, requiring a diagnosis during follow-up when the child was ≥6 years old between 1/1/2017–12/31/2023.

Covariates

Maternal socio-demographics. This includes self-reported race and ethnicity (Asian/Pacific Islander, non-Hispanic Black, Hispanic, non-Hispanic White, and Other/unknown, included as a social construct due to differences in prenatal cannabis use and child depressive and anxiety disorder prevalence by race and ethnicity), maternal age at pregnancy onset (<18, 18-24, 25-30, 31-35, 36+), parity (0, 1, 2+), insurance type (Medicaid vs. non-Medicaid), education (high school or less, some college, college graduate, and

graduate school), maternal neighborhood deprivation index (NDI; categorized into quartiles (Messer et al., 2006) and infant birth year. Maternal race/ethnicity, parity, and education were supplemented with birth certificate data if missing from the EHR.

Maternal prenatal use of other non-cannabis substances. Substance-use was assessed at entrance to prenatal care and included alcohol, nicotine, opioids, stimulants, anxiolytics/sedatives, selective serotonin reuptake inhibitors (SSRIs), and non-SSRI antidepressants (eAppendix 1).

Prenatal care initiation. The Kotelchuck Month of Initiation Index (Kotelchuck, 1994) was used, with categories for adequate plus (month 1-2), adequate (month 3-4), intermediate (month 5-6) and inadequate (month 7+).

Maternal comorbidities. Diagnoses based on ICD-9/ICD-10 were used for nausea and vomiting between pregnancy onset and first prenatal visit; asthma, thyroid disorders, chronic pain disorders, hypertension disorders in the year before pregnancy onset through first prenatal office visit; and diabetes mellitus diagnosed in the 2 years before pregnancy onset through first prenatal visit. Maternal anxiety, depressive, and other psychiatric disorders were defined using diagnoses during the year before pregnancy onset through the first prenatal visit (eAppendix 2).

Statistical Analysis

Marginal Cox proportional hazards regression models with a cluster term at the maternal level and robust standard errors to account for correlated observations (i.e., multiple pregnancies nested within individuals) were fit to estimate the association between prenatal cannabis use and outcomes. Follow-up time began when children were 72 months old (6 years), and age of child in months was used as the time scale. Children were followed until the earliest of outcome occurrence (first diagnosis), end of KPNC enrollment (>3-month gap in enrollment), no pediatric or mental health department visit within a required age window (see below), death, or end of the study period (December 31, 2023), with a maximum follow-up age of 13 years. Attendance at ≥1 pediatric or mental health visit during the intervals (age 6-9 and 9-12) was required to

ensure that clinicians had the opportunity to recognize and diagnose anxiety and depressive disorders (eAppendix 3).

Inverse probability of censoring weights (IPCW) was applied to account for the potential impact of informative censoring due to any differences in healthcare visits or membership loss among children of individuals with versus without prenatal cannabis use (Howe et al., 2016). Time-varying stabilized weights were generated for 3 censoring time intervals (months 72-107, 108-143, and 144-155).

Child anxiety and depressive disorders were modeled separately. For each outcome, a series of analyses sequentially adjusted for sets of covariates to examine the degree of confounding. Model 1 did not include any covariates. Model 2 adjusted for maternal sociodemographic characteristics, Model 3 additionally adjusted for other prenatal non-cannabis substance use, Model 4 additionally adjusted for prenatal care initiation, Model 5 additionally adjusted for maternal medical comorbidities, and Model 6 additionally adjusted for maternal mental health comorbidities. All models adjusted for child's age using age as the time scale.

For both the primary and secondary exposures, sensitivity analyses were conducted that: 1) tested for interactions by child sex on the primary exposure and outcomes, 2) excluded children born to individuals with any non-cannabis prenatal substance use, and 3) stratified by maternal anxiety disorder (for child anxiety disorder models) or by maternal depressive disorder (for child depressive disorder models),

and 4) defined prenatal cannabis use by a) only self-report and b) only urine toxicology.

Analyses were performed using SAS, version 9.4 and R, version 4.0.2, and two-sided P values <.05 were considered statistically significant.

RESULTS

Of the 115,533 pregnancies (from 97,376 unique individuals), 27.0% were Asian/Pacific Islander, 5.8% non-Hispanic Black, 24.7% Hispanic, and 38.5% were non-Hispanic White; 11.9% were aged ≤ 24 years, 4.3% were insured by Medicaid, and 15.6% were to individuals with \leq high school education (Table 1). The median gestational age at substance use screening was 8.0 (inter-quartile range [IQR]=2.6) weeks. Overall, 4.4% screened positive for cannabis use by self-report or urine toxicology testing (0.4% daily, 0.5% weekly, 1.0% monthly, and 2.5% unknown frequency); 0.7% were positive by self-report only, 2.5% were positive by urine toxicology testing only, and 1.2% were positive on both.

The cohort of children was 51.3% male, 48.7% female. The median length of follow-up was 8.8 (IQR=3.3) years. By follow-up end, 9,230 children were diagnosed with an anxiety disorder with a first diagnosis at median age 8 (IQR=1.7) years, and 1,224 children were diagnosed with a depressive disorder with median age of first diagnosis at 9.4 (IQR=1.7) years. Most were diagnosed in a mental health specialty department (anxiety disorder: 78%; depressive disorder: 93%).

Table 1. *Characteristics of 115,533 Mother-Child Dyads born 2011-2017, Overall and by Prenatal Cannabis Use*

	Pregnancies ^a , n (%)	Prenatal Cannabis Use ^b	
		No, n (%)	Yes, n (%)
Total	115,533	110,451	5,082
Maternal sociodemographic characteristics			
Race/ethnicity			
Asian/Pacific Islander	31,239 (27.0)	30,938 (28.0)	301 (5.9)
Non-Hispanic Black	6,735 (5.8)	5,530 (5.0)	1,205 (23.7)
Hispanic	28,574 (24.7)	27,233 (24.7)	1,341 (26.4)
Non-Hispanic White	44,446 (38.5)	42,545 (38.5)	1,901 (37.4)
Other ^c /Unknown	4,539 (3.9)	4,205 (3.8)	334 (6.6)
Age			
<18	969 (0.8)	783 (0.7)	186 (3.7)
18-24	12,787 (11.1)	10,939 (9.9)	1,848 (36.4)
25-30	38,002 (32.9)	36,498 (33.0)	1,504 (29.6)
31-35	42,660 (36.9)	41,541 (37.6)	1,119 (22.0)

Prenatal Cannabis and Offspring Anxiety and Depression

36+	21,115 (18.3)	20,690 (18.7)	425 (8.4)
Parity			
0	46,687 (40.4)	43,903 (39.7)	2,784 (54.8)
1	43,474 (37.6)	42,043 (38.1)	1,431 (28.2)
2+	25,372 (22.0)	24,505 (22.2)	867 (17.1)
Insurance type			
Medicaid	4,934 (4.3)	3,967 (3.6)	967 (19.0)
Non-Medicaid	110,599 (95.7)	106,484 (96.4)	4,115 (81.0)
Education level			
High School or Less	17,976 (15.6)	16,237 (14.7)	1,739 (34.2)
Some College	33,827 (29.3)	31,642 (28.6)	2,185 (43.0)
College Graduate	38,286 (33.1)	37,519 (34.0)	767 (15.1)
Graduate School	23,081 (20.0)	22,835 (20.7)	246 (4.8)
Unknown	2,363 (2.0)	2,218 (2.0)	145 (2.9)
Neighborhood Deprivation Index, Quartile			
Q1 - Least deprived	28,879 (25.0)	28,237 (25.6)	642 (12.6)
Q2	28,884 (25.0)	27,889 (25.3)	995 (19.6)
Q3	28,887 (25.0)	27,546 (24.9)	1,341 (26.4)
Q4 - Most deprived	28,883 (25.0)	26,779 (24.2)	2,104 (41.4)
Delivery year			
2011	15,164 (13.1)	14,566 (13.2)	598 (11.8)
2012	15,823 (13.7)	15,204 (13.8)	619 (12.2)
2013	16,164 (14.0)	15,512 (14.0)	652 (12.8)
2014	16,549 (14.3)	15,841 (14.3)	708 (13.9)
2015	16,627 (14.4)	15,901 (14.4)	726 (14.3)
2016	17,361 (15.0)	16,530 (15.0)	831 (16.4)
2017	17,845 (15.4)	16,897 (15.3)	948 (18.7)
Prenatal substance use^d			
Alcohol use	10,645 (9.2)	9,572 (8.7)	1,073 (21.1)
Nicotine use	4,849 (4.2)	3,515 (3.2)	1,334 (26.2)
Stimulant use	641 (0.6)	470 (0.4)	171 (3.4)
Opioid use	3,458 (3.0)	3,041 (2.8)	417 (8.2)
Anxiolytic/Sedative use	3,089 (2.7)	2,763 (2.5)	326 (6.4)
SSRI ^e use	3,803 (3.3)	3,505 (3.2)	298 (5.9)
Non-SSRI antidepressant use	1,719 (1.5)	1,545 (1.4)	174 (3.4)
Maternal prenatal care utilization			
Gestational age at prenatal care initiation ^f			
Adequate Plus (Month 1-2)	76,616 (66.3)	73,467 (66.5)	3,149 (62.0)
Adequate (Month 3-4)	37,111 (32.1)	35,371 (32.0)	1,740 (34.2)
Intermediate (Month 5-6)	1,345 (1.2)	1,209 (1.1)	136 (2.7)
Inadequate (Month 7+)	461 (0.4)	404 (0.4)	57 (1.1)
Maternal medical comorbidities			
Pre-existing asthma	11,218 (9.7)	10,345 (9.4)	873 (17.2)
Pre-existing chronic pain	3,584 (3.1)	3,242 (2.9)	342 (6.7)
Pre-existing diabetes (Type I or II)	1,641 (1.4)	1,575 (1.4)	66 (1.3)
Pre-existing hypertensive disorder	2,879 (2.5)	2,730 (2.5)	149 (2.9)
Pre-existing thyroid disorder	6,520 (5.6)	6,351 (5.8)	169 (3.3)
Nausea/vomiting in pregnancy ^g	12,147 (10.5)	10,992 (10.0)	1,155 (22.7)
Maternal mental health comorbidities			
Pre-existing anxiety disorder	3,774 (3.3)	3,486 (3.2)	288 (5.7)
Pre-existing depressive disorder	8,943 (7.7)	8,172 (7.4)	771 (15.2)
Pre-existing substance use disorder ^h	3,541 (3.1)	2,669 (2.4)	872 (17.2)
Pre-existing other psychiatric disorder	3,317 (2.9)	2,903 (2.6)	414 (8.1)
Infant characteristics			
Sex			
Female	56,337 (48.8)	53,837 (48.7)	2,500 (49.2)
Male	59,196 (51.2)	56,614 (51.3)	2,582 (50.8)

Secondary exposure

Frequency of prenatal cannabis use

None	110,451 (95.6)	110,451 (100.0)	0 (0.0)
Monthly or less	1,160 (1.0)	0 (0.0)	1,160 (22.8)
Weekly	561 (0.5)	0 (0.0)	561 (11.0)
Daily	422 (0.4)	0 (0.0)	422 (8.3)
Unknown frequency ⁱ	2,939 (2.5)	0 (0.0)	2,939 (57.8)

^aThe cohort consisted of 115,533 unique pregnancies among 97,376 unique individuals. ^bPrenatal cannabis use was determined by self-reported use and/or urine toxicology test results. ^cOther race/ethnicity includes American Indian, Alaskan-Native, and multi-racial individuals. ^dAssessed at entrance to prenatal care by self-report or urine toxicology, or determined by prescription dispensing between the date of last menstrual period and first prenatal visit, or before pregnancy with days' supply lasting past the date of last menstrual period. ^eSelective Serotonin Reuptake Inhibitor. ^fMonths gestation at first prenatal office visit. ^gDiagnosis between date of last menstrual period and first prenatal office visit. ^hExcludes cannabis-related substance use disorders. ⁱUnknown frequency includes those who did not self-report cannabis use since becoming pregnant but tested positive for THC on urine toxicology.

Anxiety disorders. Prenatal cannabis use was not associated with child anxiety disorders in Model 1 (hazard ratio [HR]:1.08, 95%CI: 0.98-1.18; Table 2). However, after adjustment for covariates, there was a statistically significant inverse association in Models 3-6 (Model 6: HR:0.84, 95%CI: 0.75-0.95; Table 2, Figure 1).

For frequency of cannabis use, in Model 1 relative to no prenatal cannabis use, monthly or less use, weekly use, and daily use were associated with higher risk of child anxiety disorders (Table 2). In Model 6, after adjusting for all covariates, relative to no use, monthly or less (HR:0.91, 95%CI: 0.73-1.14), weekly (HR:1.03, 95%CI: 0.77-1.38), and daily (HR:1.07, 95%CI: 0.76-1.51) use were not associated with offspring anxiety disorders (Table 2; Figure 1). However, unknown frequency (positive toxicology but no self-reported use) was inversely associated with child anxiety disorders (HR:0.75, 95%CI: 0.63-0.88).

Depressive disorders. Prenatal cannabis use was positively associated with child depressive disorders in Model 1 (HR:1.62, 95%CI:1.35-1.94, Table 2). After adjustment, the association was no longer significant (Model 6: HR:1.15, 95%CI: 0.85-1.54, Table 2, Figure 1).

For frequency of cannabis use, in Model 1, relative to no prenatal cannabis use, monthly or less use and unknown frequency of use were associated with higher risk of child depressive disorders (Table 2). After adjusting for covariates, monthly or less (HR:1.54, 95%CI: 0.96-2.48), weekly (HR:0.81, 95%CI: 0.36-1.83), daily (HR:0.96, 95%CI: 0.41-2.25), and unknown frequency of use (HR:1.08, 95%CI: 0.73-1.60) were not associated with child depressive disorders relative to no use (Model 6; Table 2; Figure 1).

Sensitivity Analyses

In the fully adjusted sensitivity analyses excluding individuals who screened positive for any non-cannabis prenatal substance use, prenatal cannabis use was inversely associated with child anxiety disorders (HR:0.82, 95%CI: 0.68-0.98; eTable 1) and was not associated with child depressive disorders (HR:1.00, 95%CI: 0.62-1.60; eTable 1). There were no interactions between prenatal cannabis use and child sex ($p > .05$) for child anxiety or depressive disorders. There were no associations between the frequency of prenatal cannabis use and child outcomes.

In the fully adjusted sensitivity analyses stratified by maternal anxiety or depressive disorders, results were similar for individuals with and without anxiety or depression (eTable 2).

Prenatal cannabis use was not associated with child anxiety disorders in fully adjusted models when cannabis use was defined only by self-report (HR:1.00, 95%CI: 0.85-1.18), but there was an inverse association when use was defined only by toxicology testing (HR:0.79, 95%CI: 0.69-0.90). Prenatal cannabis use was not associated with child depressive disorders when use was defined only by self-report (HR:1.20, 95%CI: 0.83-1.78) or only by a toxicology text (HR:1.00, 95%CI: 0.71-1.39).

Table 2. Hazard Ratios for Associations of Prenatal Cannabis Use and Self-reported Frequency of Use with Child Anxiety and Depression in Mother-Child Dyads Born 2011-2017

		Model 1	Model 2	Model 3	Model 4	Model 5	Model 6
	Child Anxiety, n (%)	HR (95% CI)	HR (95% CI)	HR (95% CI)	HR (95% CI)	HR (95% CI)	HR (95% CI)
Prenatal Cannabis Use							
No	8,833 (8.0)	1.00 (Ref)	1.00 (Ref)	1.00 (Ref)	1.00 (Ref)	1.00 (Ref)	1.00 (Ref)
Yes	397 (7.8)	1.08 (0.98-1.18)	0.94 (0.84-1.06)	0.86 (0.76-0.97)	0.86 (0.76-0.97)	0.85 (0.75-0.95)	0.84 (0.75-0.95)
Frequency of Cannabis Use							
None	8,830 (8.0)	1.00 (Ref)	1.00 (Ref)	1.00 (Ref)	1.00 (Ref)	1.00 (Ref)	1.00 (Ref)
Monthly or less	100 (8.6)	1.19 (1.00-1.42)	1.01 (0.81-1.26)	0.90 (0.72-1.13)	0.92 (0.73-1.15)	0.91 (0.73-1.14)	0.91 (0.73-1.14)
Weekly	57 (10.2)	1.42 (1.13-1.80)	1.19 (0.89-1.59)	1.03 (0.77-1.38)	1.03 (0.77-1.39)	1.03 (0.77-1.38)	1.03 (0.77- 1.38)
Daily	42 (10.0)	1.47 (1.11-1.93)	1.27 (0.90-1.78)	1.09 (0.77-1.54)	1.12 (0.80-1.58)	1.10 (0.78-1.55)	1.07 (0.76-1.51)
Unknown frequency ^a	198 (6.7)	0.91 (0.80-1.04)	0.82 (0.70-0.96)	0.77 (0.65-0.90)	0.76 (0.65-0.90)	0.75 (0.63-0.88)	0.75 (0.63-0.88)
	Child Depression, n (%)	HR (95% CI)	HR (95% CI)	HR (95% CI)	HR (95% CI)	HR (95% CI)	HR (95% CI)
Prenatal Cannabis Use							
No	1,153 (1.0)	1.00 (Ref)	1.00 (Ref)	1.00 (Ref)	1.00 (Ref)	1.00 (Ref)	1.00 (Ref)
Yes	71 (1.4)	1.62 (1.35-1.94)	1.34 (1.02-1.77)	1.15 (0.85-1.54)	1.15 (0.86-1.54)	1.15 (0.85-1.54)	1.15 (0.85-1.54)
Frequency of Cannabis Use							
None	1,153 (1.0)	1.00 (Ref)	1.00 (Ref)	1.00 (Ref)	1.00 (Ref)	1.00 (Ref)	1.00 (Ref)
Monthly or less	23 (2.0)	2.18 (1.58-3.01)	1.86., (1.17-2.95)	1.50 (0.93-2.42)	1.53 (0.95-2.46)	1.53 (0.95-2.45)	1.54 (0.96-2.48)
Weekly	7 (1.2)	1.38 (0.78-2.43)	1.07 (0.48, 2.38)	0.80 (0.36-1.80)	0.80 (0.36-1.80)	0.81 (0.36-1.81)	0.81 (0.36-1.83)
Daily	7 (1.7)	1.77 (0.95-3.29)	1.29 (0.55, 3.02)	0.98 (0.42-2.29)	0.99 (0.42-2.32)	0.99 (0.42-2.32)	0.96 (0.41-2.25)
Unknown frequency ^a	34 (1.2)	1.42 (1.10-1.83)	1.20 (0.82, 1.75)	1.09 (0.74-1.61)	1.09 (0.74-1.61)	1.09 (0.74-1.61)	1.08 (0.73-1.60)

^a Unknown frequency includes those who did not self-report cannabis use since becoming pregnant but tested positive for THC on urine toxicology.

Model 1: Cox proportional hazards model with child age (in months) as time scales, with inverse-probability of censoring weights (IPCW) applied and maternal-level cluster term, and no additional covariates.

Model 2: Adjusted for maternal demographics (age at pregnancy onset, race/ethnicity, education, neighborhood deprivation index, parity, Medicaid insurance, delivery year).


Model 3: Additionally adjusted for other non-cannabis substance exposure (alcohol, nicotine, opioids, stimulants, anxiolytics/sedatives, SSRIs, non-SSRI antidepressants).

Model 4: Additionally adjusted for month of prenatal care initiation.

Model 5: Additionally adjusted for maternal medical comorbidities (asthma, diabetes mellitus, nausea/vomiting during pregnancy, chronic pain, thyroid disorders, hypertensive disorders).

Model 6: Additionally adjusted for maternal mental health comorbidities (anxiety disorders, depression, substance use disorders, other psychiatric disorders)

Figure 1. Adjusted Hazard Ratios for Associations of Maternal Prenatal Cannabis Use with Child Anxiety and Depression



DISCUSSION

In this large, retrospective cohort study of maternal-child dyads in a diverse, integrated healthcare delivery system, prenatal cannabis use was not associated with increased risk of child early-onset anxiety or depressive disorders after adjusting for key covariates, including maternal use of other non-cannabis substances during early pregnancy and maternal pre-pregnancy anxiety and depressive disorders.

In unadjusted analyses, prenatal cannabis use was associated with increased risk of early-onset depressive disorders and was not associated with early-onset anxiety disorders; however, the association with depressive disorders was attenuated after sequential adjustment for maternal sociodemographic characteristics, other

prenatal substance use, and maternal comorbidities. These findings suggest that the initial unadjusted associations may reflect common underlying risk factors rather than an independent effect of prenatal cannabis use on offspring depressive disorders.

Several small studies have evaluated the relationship between prenatal cannabis use and child anxiety outcomes from ages 2-9, using parent-reported screening measures or structured interviews (Moore et al., 2023; Nomura et al., 2023; O'Connell & Fried, 1991; Rompala et al., 2021; Sullivan et al., 2024). The number of pregnancies with cannabis use ranged from 6 to 68, and findings were mixed, with only one study reporting an increased risk of child anxiety associated with prenatal cannabis use (Rompala et al., 2021), while the remaining four found no

association (Moore et al., 2023; Nomura et al., 2023; O'Connell & Fried, 1991; Sullivan et al., 2024). There are several potential explanations for the lower risk of offspring anxiety disorders associated with prenatal cannabis use in our study. For example, individuals who use cannabis during pregnancy may be different in other key ways that could contribute to lower risk of offspring anxiety disorders (e.g., different lifestyle behaviors, support networks, or parenting approaches). A supportive postnatal environment can potentially impact offspring brain development, including cognitive ability, cell proliferation, and synaptic protein expression, induced by adverse environmental exposures in utero (Koo et al., 2003; Soeda et al., 2010). Further, those who use cannabis could be less likely to recognize symptoms of anxiety in their children or less likely to seek medical care for anxiety symptoms. Importantly, having a positive toxicology test for prenatal cannabis use but not disclosing use by self-report was associated with lower risk of a child anxiety disorder diagnosis, and those who do not report their cannabis use may also underreport anxiety symptoms in their children.

Maternal stress and depression during pregnancy and postpartum have also been shown to play a large role in offspring development and mental health (Rogers et al., 2020). Pregnant individuals report using cannabis for stress and anxiety relief (Skelton et al., 2020; Vanstone et al., 2021), and there is emerging evidence suggesting the potential therapeutic benefit of cannabis on depression symptoms, as an anxiolytic, for stress relief, and improved sleep quality (Ayisire et al., 2022). However, meta-analyses suggest that long-term cannabis use is associated with worse mental health outcomes, particularly in vulnerable populations (Lacasse et al., 2023; Xue et al., 2021). Maternal cannabis use might have a protective effect on offspring anxiety disorders by mitigating maternal stress and other psychosocial factors impacting offspring development, highlighting the importance of early interventions to support those transitioning into parenthood with mental health resources.

Our finding of no association between prenatal cannabis use and early onset of offspring depressive disorders is consistent with two of three smaller studies investigating the

association between prenatal cannabis use, assessed by self-report (Gray et al., 2005; Rompala et al., 2021) or urine toxicology testing (Moore et al., 2023), and child depression symptoms from ages 3-10, with the number of pregnancies with prenatal cannabis use ranging from 6 to 268. Notably, an older study of pregnancies from 1982-1985, reported a positive association between self-reported prenatal cannabis use and child depression (Gray et al., 2005). However, a more recent study of pregnancies from 2010-2014, relying on urine toxicology testing in the second trimester, found no association with depressive or anxiety problems based on the DSM-oriented Child Behavior Checklist (CBCL) and an inverse association with mean internalizing behaviors (Moore et al., 2023). Additionally, a study of pregnancies from 2010-2015 (Rompala et al., 2021) found that self-reported prenatal cannabis use was not associated with child depression on the Behavior Assessment System for Children, Second Edition (BASC-2).

Strengths and Limitations

This study has several strengths, including the large and diverse sample of pregnant individuals who were universally screened for prenatal cannabis use by self-report and urine toxicology testing during standard prenatal care. KPNC has universal screening for parent-reported child depression symptoms as part of standard well child visits, increasing the likelihood that children with symptoms were identified, assessed, and given a diagnosis when needed. Further, the cohort included children born between 2011 and 2017, allowing us to examine outcomes in the context of higher strength cannabis products that have become prevalent in recent years. We also adjusted for a range of potential confounders assessed prior to or concurrent with prenatal cannabis screening and conducted sensitivity analyses that excluded individuals with any non-cannabis prenatal substance use. We did not adjust for post-exposure or downstream variables that could be on the causal pathway from cannabis to offspring mental health outcomes, and consistent findings across sensitivity analyses support the interpretation that attenuation in adjusted results reflects confounding rather than over-adjustment.

Finally, the number of pregnancies with prenatal cannabis use in our sample was nearly twenty times larger than the largest study to date examining prenatal cannabis use and child depression or anxiety.

Our study also has limitations. The sample was limited to insured pregnant individuals in California, where cannabis has been legal for medical use since 1996 and for adult recreational use since 2016, and results may not generalize to uninsured individuals or to those in other states. Further, our cohort was limited to pregnant individuals who were enrolled with KPNC in the year prior to pregnancy through delivery, and to children who were enrolled with KPNC at age 6. These exclusions contributed to selection bias, as pregnant individuals who entered prenatal care later were more likely to use cannabis, and children who left KPNC may be at a greater risk for adverse outcomes. Additionally, while prenatal cannabis use was universally assessed by self-report and toxicology testing as part of standard care, we cannot distinguish between use in pregnancy that stopped after recognition of pregnancy versus use that continued throughout pregnancy. We did not have data on paternal cannabis use, mode of use, or secondhand cannabis exposure, which may impact offspring outcomes. Although we adjusted for a range of covariates in the EHR, unmeasured confounding cannot be ruled out. However, we applied inverse probability of censoring weights calculated using maternal sociodemographic characteristics, prenatal substance use, pre-existing medical comorbidities, and prenatal care utilization to account for differential healthcare utilization patterns, reducing the impact of informative censoring on model results. Still, applying the IPCW may have not fully eliminated informative censoring bias due to unmeasured factors not included in the weights. In addition, the IPCW did not address potential selection bias introduced by child loss to follow-up before age 6 (Avalos et al., 2023). While we followed children up to a maximum age of 13 years, our results reflect early onset of depressive and anxiety disorders, and future research with longer follow-up is needed as some may go on to develop these disorders. Finally, given the importance of social determinants of health in shaping the risk for and development of mental health problems among children and adolescents, it is crucial that future

studies examine whether associations between prenatal cannabis use and offspring mental health vary by key social determinants of health.

Conclusions

In a large, diverse cohort of pregnant individuals and their children, maternal cannabis use during early pregnancy was not associated with an increased risk of child anxiety or depressive disorders. Additional research is needed to test whether the association between prenatal cannabis use and child anxiety and depressive disorders varies depending on the timing, duration, mode and strength of cannabis. Despite the findings of no increased risk for these mental health outcomes in this study, prenatal cannabis use remains associated with adverse neonatal and maternal outcomes and national guidelines recommend discontinuation of cannabis use during pregnancy (Committee on Obstetric Practice, 2017).

REFERENCES

- Avalos, L. A., Adams, S. R., Alexeeff, S. E., Oberman, N. R., Does, M. B., Ansley, D., Goler, N., Padon, A. A., Silver, L. D., & Young-Wolff, K. C. (2024). Neonatal outcomes associated with in utero cannabis exposure: A population-based retrospective cohort study. *American Journal of Obstetrics & Gynecology*, *231*(1), 132 e131–132 e113. <https://doi.org/https://doi.org/10.1016/j.ajog.2023.11.1232>
- Avalos, L. A., Oberman, N., Alexeeff, S. E., Croen, L. A., Adams, S. R., Davignon, M., & Young-Wolff, K. C. (2023). Association between maternal prenatal cannabis use and missed child preventive care visits in an integrated health care delivery system in Northern California. *Preventative Medicine*, *175*, 107716. <https://doi.org/10.1016/j.ympmed.2023.107716>
- Avalos, L. A., Oberman, N., Alexeeff, S. E., Croen, L. A., Davignon, M. N., Adams, S. R., Ansley, D., Chambers, C. D., Steuerle, K., & Young-Wolff, K. C. (2024). Early maternal prenatal cannabis use and child developmental delays. *JAMA Network Open*, *7*(10), e2440295.

- <https://doi.org/10.1001/jamanetworkopen.2024.40295>
- Avalos, L. A., Shenkute, M., Alexeeff, S. E., Oberman, N., Croen, L. A., Davignon, M., Adams, S. R., Ansley, D., Castellanos, C., & Young-Wolff, K. C. (2024). Maternal prenatal cannabis use and child autism spectrum disorder. *JAMA Network Open*, 7(10), e2440301. <https://doi.org/10.1001/jamanetworkopen.2024.40301>
- Ayisire, O. E., Okobi, O. E., Adaralegbe, N. J., Adeosun, A. A., Sood, D., Onyechi, N. P., Agazie, O., Shittu, H. O., Akinsola, Z., Nnaji, C. G., Owolabi, O. J., Umeh, N. J., Imobighe, I. C., Adedoyin, A. M., & Usman, M. (2022). The use of cannabis and its effects on postpartum depression. *Cureus*, 14(8), e27926. <https://doi.org/https://doi.org/10.7759/cureus.27926>
- Baia, I., & Domingues, R. M. S. M. (2022). The effects of cannabis use during pregnancy on low birth weight and preterm birth: A systematic review and meta-analysis. *American Journal of Perinatology*. <https://doi.org/https://doi.org/10.1055/a-1911-3326>
- Bandoli, G., Jelliffe-Pawłowski, L., Schumacher, B., Baer, R. J., Felder, J. N., Fuchs, J. D., Oltman, S. P., Steurer, M. A., & Marienfeld, C. (2021). Cannabis-related diagnosis in pregnancy and adverse maternal and infant outcomes. *Drug and Alcohol Dependence*, 225, 108757. <https://doi.org/10.1016/j.drugalcdep.2021.108757>
- Bassalov, H., Yakirevich-Amir, N., Reuveni, I., Monk, C., Florentin, S., Bonne, O., & Matok, I. (2024). Prenatal cannabis exposure and the risk for neuropsychiatric anomalies in the offspring: a systematic review and meta-analysis. *American Journal of Obstetrics & Gynecology*. <https://doi.org/10.1016/j.ajog.2024.06.014>
- Bayrampour, H., Zahradnik, M., Lisonkova, S., & Janssen, P. (2019). Women's perspectives about cannabis use during pregnancy and the postpartum period: An integrative review. *Preventative Medicine*, 119, 17–23. <https://doi.org/10.1016/j.ypmed.2018.12.002>
- Blackard, C., & Tennes, K. (1984). Human placental transfer of cannabinoids. *New England Journal of Medicine*, 311(12), 797. <https://doi.org/10.1056/NEJM198409203111213>
- Chabbarria, K. C., Racusin, D. A., Antony, K. M., Kahr, M., Suter, M. A., Mastrobattista, J. M., & Aagaard, K. M. (2016). Marijuana use and its effects in pregnancy. *American Journal of Obstetrics & Gynecology*, 215(4), 506 e501–507. <https://doi.org/10.1016/j.ajog.2016.05.044>
- Chang, J. C., Tarr, J. A., Holland, C. L., De Genna, N. M., Richardson, G. A., Rodriguez, K. L., Sheeder, J., Kraemer, K. L., Day, N. L., Rubio, D., Jarlenski, M., & Arnold, R. M. (2019). Beliefs and attitudes regarding prenatal marijuana use: Perspectives of pregnant women who report use. *Drug and Alcohol Dependence*, 196, 14–20. <https://doi.org/https://doi.org/10.1016/j.drugalcdep.2018.11.028>
- Cioffredi, L. A., Anderson, H., Loso, H., East, J., Nguyen, P., Garavan, H., & Potter, A. (2022). Prenatal cannabis exposure predicts attention problems, without changes on fMRI in adolescents. *Neurotoxicology and Teratology*, 91, 107089. <https://doi.org/10.1016/j.ntt.2022.107089>
- Committee on Obstetric Practice. (2017). *Committee Opinion No. 722: Marijuana use during pregnancy and lactation*. (2017). *Obstetrics & Gynecology*, 130(4), e205–e209. <https://doi.org/10.1097/AOG.0000000000000235>
- Corsi, D. J., Donelle, J., Sucha, E., Hawken, S., Hsu, H., El-Chaâr, D., Bisnaire, L., Fell, D., Wen, S. W., & Walker, M. (2020). Maternal cannabis use in pregnancy and child neurodevelopmental outcomes. *Nature Medicine*, 26(10), 1536–1540. <https://doi.org/10.1038/s41591-020-1002-5>
- Corsi, D. J., Walsh, L., Weiss, D., Hsu, H., El-Chaar, D., Hawken, S., Fell, D. B., & Walker, M. (2019). Association between self-reported prenatal cannabis use and maternal, perinatal, and neonatal outcomes. *JAMA*, 322(2), 145–152. <https://doi.org/10.1001/jama.2019.8734>

- De Genna, N. M., Willford, J. A., & Richardson, G. A. (2022). Long-term effects of prenatal cannabis exposure: pathways to adolescent and adult outcomes. *Pharmacology Biochemistry and Behavior*, *214*, 173358. <https://doi.org/10.1016/j.pbb.2022.173358>
- DISA Global Solutions. *Marijuana legality by state — Updated September 2, 2024*. <https://disa.com/marijuana-legality-by-state>
- ElSohly, M. A., Mehmedic, Z., Foster, S., Gon, C., Chandra, S., & Church, J. C. (2016). Changes in cannabis potency over the last 2 decades (1995-2014): analysis of current data in the United States. *Biological Psychiatry*, *79*(7), 613–619. <https://doi.org/10.1016/j.biopsych.2016.01.004>
- Foti, T. R., Green, A., Altschuler, A., Iturralde, E., Does, M. B., Jackson-Morris, M., Adams, S. R., Goler, N., Ansley, D., Conway, A., & Young-Wolff, K. C. (2023). Patient perceptions of prenatal cannabis use and implications for clinicians. *Obstetrics & Gynecology*, *142*(5), 1153–1161. <https://doi.org/https://doi.org/10.1097/AOG.0000000000005295>
- Fried, P. A., & Watkinson, B. (1988). 12- and 24-month neurobehavioural follow-up of children prenatally exposed to marihuana, cigarettes and alcohol. *Neurotoxicology and Teratology*, *10*(4), 305–313. [https://doi.org/10.1016/0892-0362\(88\)90032-3](https://doi.org/10.1016/0892-0362(88)90032-3)
- Fried, P. A., & Watkinson, B. (1990). 36- and 48-month neurobehavioral follow-up of children prenatally exposed to marijuana, cigarettes, and alcohol. *Journal of Developmental & Behavioral Pediatrics*, *11*(2), 49–58. <https://www.ncbi.nlm.nih.gov/pubmed/2324288>
- Fried, P. A., Watkinson, B., & Gray, R. (1992). A follow-up study of attentional behavior in 6-year-old children exposed prenatally to marihuana, cigarettes, and alcohol. *Neurotoxicology and Teratology*, *14*(5), 299–311. [https://doi.org/10.1016/0892-0362\(92\)90036-a](https://doi.org/10.1016/0892-0362(92)90036-a)
- Garrison-Desany, H. M., Hong, X., Maher, B. S., Beaty, T. H., Wang, G., Pearson, C., Liang, L., Wang, X., & Ladd-Acosta, C. (2022). Individual and combined association between prenatal polysubstance exposure and childhood risk of attention-deficit/hyperactivity disorder. *JAMA Network Open*, *5*(3), e221957. <https://doi.org/10.1001/jamanetworkopen.2022.1957>
- Grant, K. S., Conover, E., & Chambers, C. D. (2020). Update on the developmental consequences of cannabis use during pregnancy and lactation. *Birth Defects Research*, *112*(15), 1126–1138. <https://doi.org/10.1002/bdr2.1766>
- Gray, K. A., Day, N. L., Leech, S., & Richardson, G. A. (2005). Prenatal marijuana exposure: Effect on child depressive symptoms at ten years of age. *Neurotoxicology and Teratology*, *27*(3), 439–448. <https://doi.org/10.1016/j.ntt.2005.03.010>
- Howe, C. J., Cole, S. R., Lau, B., Napravnik, S., & Eron, J. J., Jr. (2016). Selection bias due to loss to follow up in cohort studies. *Epidemiology*, *27*(1), 91–97. <https://doi.org/10.1097/EDE.0000000000000409>
- Koo, J. W., Park, C. H., Choi, S. H., Kim, N. J., Kim, H. S., Choe, J. C., & Suh, Y. H. (2003). The postnatal environment can counteract prenatal effects on cognitive ability, cell proliferation, and synaptic protein expression. *FASEB Journal*, *17*(11), 1556–1558. <https://doi.org/10.1096/fj.02-1032fje>
- Kotelchuck, M. (1994). An evaluation of the Kessner Adequacy of Prenatal Care Index and a proposed Adequacy of Prenatal Care Utilization Index. *American Journal of Public Health*, *84*(9), 1414–1420. <https://doi.org/https://doi.org/10.2105/ajph.84.9.1414>
- Koto, P., Allen, V. M., Fahey, J., & Kuhle, S. (2022). Maternal cannabis use during pregnancy and maternal and neonatal outcomes: A retrospective cohort study. *BJOG*, *129*(10), 1687–1694. <https://doi.org/10.1111/1471-0528.17114>
- Lacasse, A., Nguena Nguefack, H. L., Page, G., Choiniere, M., Samb, O. M., Katz, J., Menard, N., Vissandjee, B., Zerriouh, M., & TorSa, D. E. C. W. (2023). Sex and gender differences in healthcare utilisation trajectories: A cohort study among Quebec workers living with chronic pain. *BMJ Open*, *13*(7), e070509. <https://doi.org/https://doi.org/10.1136/bmjopen-2022-070509>
- Lee, C. C., & Chiang, C. N. (1985). Maternal-fetal transfer of abused substances:

- pharmacokinetic and pharmacodynamic data. *NIDA Research Monograph*, 60, 110–147. <https://www.ncbi.nlm.nih.gov/pubmed/2997613>
- Lo, J. O., Shaw, B., Robalino, S., Ayers, C. K., Durbin, S., Rushkin, M. C., Olyaei, A., Kansagara, D., & Harrod, C. S. (2023). Cannabis use in pregnancy and neonatal outcomes: A systematic review and meta-analysis. *Cannabis and Cannabinoid Research*. <https://doi.org/10.1089/can.2022.0262>
- Messer, L. C., Laraia, B. A., Kaufman, J. S., Eyster, J., Holzman, C., Culhane, J., Elo, I., Burke, J. G., & O'Campo, P. (2006). The development of a standardized neighborhood deprivation index. *Journal of Urban Health*, 83(6), 1041–1062. <https://doi.org/10.1007/s11524-006-9094-x>
- Moore, B. F., Salmons, K. A., Hoyt, A. T., Swenson, K. S., Bates, E. A., Sauder, K. A., Shapiro, A. L. B., Wilkening, G., Kinney, G. L., Neophytou, A. M., Sempio, C., Klawitter, J., Christians, U., & Dabelea, D. (2023). Associations between prenatal and postnatal exposure to cannabis with cognition and behavior at age 5 years: The Healthy Start Study. *International Journal of Environmental Research and Public Health*, 20(6). <https://doi.org/10.3390/ijerph20064880>
- Mulligan, M. K., & Hamre, K. M. (2023). Influence of prenatal cannabinoid exposure on early development and beyond. *Advances in Drug and Alcohol Research*, 3(4), 10981. <https://doi.org/10.3389/adar.2023.10981>
- Nomura, Y., Ham, J., Pehme, P. M., Wong, W., Pritchett, L., Rabinowitz, S., Foldi, N. S., Hinton, V. J., Wickramaratne, P. J., & Hurd, Y. L. (2023). Association of maternal exposure to Superstorm Sandy and maternal cannabis use with development of psychopathology among offspring: the Stress in Pregnancy Study. *BJPsych Open*, 9(3), e94. <https://doi.org/10.1192/bjo.2022.595>
- O'Connell, C. M., & Fried, P. A. (1991). Prenatal exposure to cannabis: a preliminary report of postnatal consequences in school-age children. *Neurotoxicology and Teratology*, 13(6), 631–639. [https://doi.org/10.1016/0892-0362\(91\)90047-z](https://doi.org/10.1016/0892-0362(91)90047-z)
- Paul, S. E., Hatoum, A. S., Fine, J. D., Johnson, E. C., Hansen, I., Karcher, N. R., Moreau, A. L., Bondy, E., Qu, Y., Carter, E. B., Rogers, C. E., Agrawal, A., Barch, D. M., & Bogdan, R. (2021). Associations between prenatal cannabis exposure and childhood outcomes: Results from the ABCD Study. *JAMA Psychiatry*, 78(1), 64–76. <https://doi.org/10.1001/jamapsychiatry.2020.2902>
- Paul, S. E., Rogers, C. E., & Bogdan, R. (2023). Pre-natal cannabis exposure: associations with development and behaviour. In D. C. D'Souza, D. J. Castle, & R. M. Murray (Eds.), *Marijuana and Madness* (3rd ed., pp. 267–278). Cambridge University Press. <https://www.cambridge.org/us/universitypress/subjects/medicine/mental-health-psychiatry-and-clinical-psychology/marijuana-and-madness-3rd-edition?format=PB&isbn=9781009305433>
- Petrangelo, A., Czuzoj-Shulman, N., Balayla, J., & Abenheim, H. A. (2019). Cannabis abuse or dependence during pregnancy: A population-based cohort study on 12 million births. *Journal of Obstetrics and Gynaecology Canada*, 41(5), 623–630. <https://doi.org/10.1016/j.jogc.2018.09.009>
- Rogers, A., Obst, S., Teague, S. J., Rossen, L., Spry, E. A., Macdonald, J. A., Sunderland, M., Olsson, C. A., Youssef, G., & Hutchinson, D. (2020). Association between maternal perinatal depression and anxiety and child and adolescent development: A meta-analysis. *JAMA Pediatrics*, 174(11), 1082–1092. <https://doi.org/10.1001/jamapediatrics.2020.2910>
- Rompala, G., Nomura, Y., & Hurd, Y. L. (2021). Maternal cannabis use is associated with suppression of immune gene networks in placenta and increased anxiety phenotypes in offspring. *Proceedings of the National Academy of Science USA*, 118(47). <https://doi.org/10.1073/pnas.2106115118>
- Sharapova, S. R., Phillips, E., Sirocco, K., Kaminski, J. W., Leeb, R. T., & Rolle, I. (2018). Effects of prenatal marijuana exposure on neuropsychological outcomes in children aged 1-11 years: A systematic review. *Paediatric and Perinatal*

- Epidemiology*, 32(6), 512–532.
<https://doi.org/10.1111/ppe.12505>
- Skelton, K. R., Hecht, A. A., & Benjamin-Neelon, S. E. (2020). Women's cannabis use before, during, and after pregnancy in New Hampshire. *Preventative Medicine Reports*, 20, 101262.
<https://doi.org/10.1016/j.pmedr.2020.101262>
- Soeda, F., Tanaka, A., Shirasaki, T., & Takahama, K. (2010). An enriched environment mitigates the brain-disruptive effects of prenatal diethylstilbestrol exposure in mice. *Neuroscience*, 169(1), 223–228.
<https://doi.org/10.1016/j.neuroscience.2010.04.047>
- Sujan, A. C., Young-Wolff, K. C., & Avalos, L. A. (2022). In-utero cannabis exposure and long-term psychiatric and neurodevelopmental outcomes: the limitations of existing literature and recommendations for future research. *Birth Defects Research*, 114(13), 689–713. <https://doi.org/10.1002/bdr2.2060>
- Sullivan, E. V., Zahr, N. M., Zhao, Q., Pohl, K. M., Sassoon, S. A., & Pfefferbaum, A. (2024). Contributions of cerebral white matter hyperintensities to postural instability in aging with and without alcohol use disorder. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, 9(10), 998–1009.
<https://doi.org/10.1016/j.bpsc.2024.03.005>
- Tchuente, V., Sheehy, O., Zhao, J. P., Gorgui, J., Gomez, Y. H., & Berard, A. (2022). Is in-utero exposure to cannabis associated with the risk of attention deficit with or without hyperactivity disorder? A cohort study within the Quebec Pregnancy Cohort. *BMJ Open*, 12(8), e052220.
<https://doi.org/10.1136/bmjopen-2021-052220>
- Torres, C. A., Medina-Kirchner, C., O'Malley, K. Y., & Hart, C. L. (2020). Response: Commentary: Totality of the evidence suggests prenatal cannabis exposure does not lead to cognitive impairments: A systematic and critical review. *Frontiers in Psychology*, 11, 583516.
<https://doi.org/10.3389/fpsyg.2020.583516>
- Vanstone, M., Taneja, S., Popoola, A., Panday, J., Greyson, D., Lennox, R., & McDonald, S. D. (2021). Reasons for cannabis use during pregnancy and lactation: A qualitative study. *Cmaj*, 193(50), E1906–E1914.
<https://doi.org/10.1503/cmaj.211236>
- Volkow, N. D., Han, B., Compton, W. M., & McCance-Katz, E. F. (2019). Self-reported medical and nonmedical cannabis use among pregnant women in the United States. *JAMA*, 322(2), 167–169.
<https://doi.org/10.1001/jama.2019.7982>
- Warshak, C. R., Regan, J., Moore, B., Magner, K., Kritzer, S., & Van Hook, J. (2015). Association between marijuana use and adverse obstetrical and neonatal outcomes. *Journal of Perinatology*, 35(12), 991–995.
<https://doi.org/10.1038/jp.2015.120>
- Xue, S., Husain, M. I., Zhao, H., & Ravindran, A. V. (2021). Cannabis Use and Prospective Long-Term Association with Anxiety: A Systematic Review and Meta-Analysis of Longitudinal Studies: Usage du cannabis et association prospective à long terme avec l'anxiété: une revue systématique et une méta-analyse d'études longitudinales. *Canadian Journal of Psychiatry*, 66(2), 126–138.
<https://doi.org/10.1177/0706743720952251>
- Young-Wolff, K. C., Adams, S. R., Alexeeff, S. E., Zhu, Y., Chojolan, E., Slama, N. E., Does, M. B., Silver, L. D., Ansley, D., Castellanos, C. L., & Avalos, L. A. (2024). Prenatal cannabis use and maternal pregnancy outcomes. *JAMA Internal Medicine*, 184(9), 1083–1093.
<https://doi.org/https://doi.org/10.1001/jamainternmed.2024.3270>
- Young-Wolff, K. C., Chi, F. W., Lapham, G. T., Alexeeff, S. E., Does, M. B., Ansley, D., & Campbell, C. I. (2024). Changes in prenatal cannabis use among pregnant individuals from 2012 to 2022. *Obstetrics & Gynecology*, 144(4), e101–e104.
<https://doi.org/10.1097/AOG.00000000000005711>
- Young-Wolff, K. C., Foti, T. R., Green, A., Altschuler, A., Does, M. B., Jackson-Morris, M., Adams, S. R., Ansley, D., Conway, A., Goler, N., Mian, M. N., & Iturralde, E. (2022). Perceptions about cannabis following legalization among pregnant individuals with prenatal cannabis use in California. *JAMA Network Open*, 5(12), e2246912.
<https://doi.org/10.1001/jamanetworkopen.2022.46912>
- Young-Wolff, K. C., Kong, K., Alexeeff, S. E., Croen, L. A., Oberman, N., Kirane, H., Ansley, D., Davignon, M., Adams, S. R., &

Avalos, L. A. (2025). Prenatal cannabis use and offspring attention deficit hyperactivity disorder and disruptive behavior disorders: a retrospective cohort study. *Journal of Developmental and Behavioral Pediatrics*, 46(1), e25–e32.

<https://doi.org/10.1097/DBP.0000000000001323>

Young-Wolff, K. C., Sarovar, V., Alexeeff, S. E., Adams, S. R., Tucker, L. Y., Conway, A., Ansley, D., Goler, N., Armstrong, M. A., & Weisner, C. (2020). Trends and correlates of self-reported alcohol and nicotine use among women before and during pregnancy, 2009–2017. *Drug and Alcohol Dependence*, 214, 108168.

<https://doi.org/10.1016/j.drugalcdep.2020.108168>

Young-Wolff, K. C., Tucker, L. Y., Alexeeff, S., Armstrong, M. A., Conway, A., Weisner, C., & Goler, N. (2017). Trends in self-reported and biochemically tested marijuana use among pregnant females in California from 2009–2016 [Research Letter]. *JAMA Psychiatry*, 318(24), 2490–2491.

<https://doi.org/10.1001/jama.2017.17225>

Funding and Acknowledgements: This study was supported by grants R01DA047405 and K01DA043604 funded by National Institute on Drug Abuse (NIDA) and grant R01DA048033 co-funded by NIDA and Office of the Director, NIH (OD). The authors have no conflicts of interest to disclose.

Authorship Contribution Statement:

Dr. Kelly Young-Wolff received funding, conceptualized the study, assisted with interpretation of data, drafted the manuscript, and provided critical revisions to the manuscript.

Mahlet Shenkute and Nina Oberman extracted and analyzed the data from the electronic health record, assisted with interpretation of data, drafted the manuscript, and provided critical revisions to the manuscript.

Dr. Stacey E. Alexeeff provided important biostatistical support, assisted with interpretation of data, and critical feedback and revisions on the manuscript.

Drs. Lisa A. Croen, Jamie O. Lo, Deborah Ansley, Carley Castellanos, Kathryn K. Ridout, and Matthew E. Hirschtritt assisted with interpretation of data and provided critical feedback and revisions to the manuscript.

Dr. Lyndsay A. Avalos received funding, conceptualized the study, assisted with interpretation of data and provided critical revisions to the manuscript.

All authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

Copyright: © 2026 Authors et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by-nc-nd/4.0/), which permits unrestricted use, distribution, and reproduction, provided the original author and source are credited, the original sources is not modified, and the source is not used for commercial purposes.

Early View Date: March 13, 2026

